Seasonal Climate Forecasting Applied to Wildland Fire Management in Alaska

Uma S. Bhatt, A Sampath, PA Bieniek, A York, R Ziel, B Brettschneider, R
Thoman, H Strader, S Alden, R Jandt, G Branson, P Peng
NGGPS/MAPP PI Meeting Aug 2-3, 2017, College Park, Maryland
Thursday August 3, 2017




Aggie Creek Fire (Summer 2015), 30 miles north of Fairbanks, US Forest Service photograph

 Goal: Use seasonal forecasts to make products for decision support several months before fire season.

NOAA Grant #: NA16OAR4310142

- Boreal wildland fire in Alaska burns many acres.
- Fires are costly (e.g.,
 The record fire year of 2004 resulted in 6.5 million acres burned and was costly from property loss (> \$35M) and emergency personnel (> \$17M).
- Information a season in advance (March) would help decision makers and has potential to save \$ by allocating resources more effectively.


Motivation



Acres burned in Alaska 2017 0.6 Million acres

[Updated from Partain et al. 2016]

Three items are needed for wildland fire






Data

- Apply NMME (Kirtman et al. 2014) seasonal forecasts to wildland fire products in Alaska.
- 'Observations' Dynamically downscaled ERA-Interim for Alaska . [Bieniek et al. 2016]
- Calculate Canadian Forest Fire Weather Indices.
- Most acres burned are remote and lightning provides ignition. Lightning data is collected by BLM.


Dynamical Downscaling serves as 'Observations'



- WRF has more detailed terrain
- Reanalysis has broader high elevation areas
- Better resolves mesoscale features
- Downscaled temperature and precipitation more realistic

Canadian Forest Fire Weather Index System


Fire fighter field-use table




Panel displays categories of fire danger for spruce (http://mesowest.org/akff/captures/InteriorSpruce.png) based on the indices.

- Data needed: early afternoon values of
 - •2-m air temperature
 - relative humidity
 - 10-m winds
 - daily total precipitation.
- Expand currently available
 48 hour lead CFFWIS to
 longer leads.

Canadian Forest Fire Weather Index System




Buildup Index Climatology from 'Obs' & CFSv2 forecast




Summer Buildup Index is underestimated


'Observed' Interior AK average BUI




Station-based BUI: Large magnitudes




'Obs' larger & seasonal shift compared to CFSv2 BUI




CFSv2 BUI for Interior Alaska for record




Climatological 2-m Temperature



Climatological Precipitation




Climatological First Snow-Free Day




M. BuKader & R. Ziel


Use station data for correcting 'Obs' and then correcting forecasts.

2017 Fire Season in Alaska, small acreage burned




2017 March Forecast Average Interior BUI






Lightning strike map from last half of July 2017





Lightning observations & 'Obs' convective precip



Can we use seasonal forecasts to anticipate the probability of lightning?


- through convective precipitation & SOMs





Take Home Message

- Bias correction needed for Indices to be useful to fire mangers (Quantile mapping).
- Multidisciplinary team needed to co-produce useful products.



Acknowledgements: This work was made possible through financial support from NOAA grant NA16OAR4310142.

Thank you for your attention!



The boundary organization Alaska Fire Science Consortium is critical for this research!